

# **pPb cross-sections at LHC and high energy pp, pA and eA collisions in DIPSY**

**G. Gustafson <sup>1</sup> L. Lönnblad <sup>1</sup> , A. Ster <sup>2</sup> and T. Csörgő <sup>2,3</sup>**

**<sup>1</sup> University of Lund, Lund, Sweden**

**<sup>2</sup> Wigner Research Center for Physics, Budapest, Hungary**

**<sup>3</sup> KRF, Gyöngyös, Hungary**

**Motivation:**

**pPb@LHC and dAu@RHIC**

**The Lund Dipole Cascade Model DIPSY**

**New: nuclei in DIPSY**

**Monte Carlo cross section results**

**Comparison with Glauber results**

**(quasi-)elastic pPb cross-section at 8 TeV**

**[arXiv:1103.4320](#)**

**[arXiv:1103.4321](#)**

**[arXiv:1206.1733](#)**

**+ [arXiv:1506.09095](#) +**

# Introduction: Glauber theory++

$$p \propto \exp\left(-\sigma_{in}^{hN} \rho L\right)$$

$$T_A(b) = \int \rho(z, b) dz$$

$$\int d^2b T_A(b) = A$$

$$T_{AB}(b) = \int d^2s T_A(s) T_B(|s - b|)$$

$$\sigma_{AB} = \int d^2b \int d^2s_1^A \dots d^2s_A^A d^2s_1^B \dots d^2s_B^B \times$$

$$T_A(s_1^A) \dots T_A(s_A^A) T_B(s_1^B) \dots T_B(s_B^B) \times$$

$$\left\{ 1 - \prod_{j=1}^B \prod_{i=1}^A \left[ 1 - \sigma(b - s_i^A + s_j^B) \right] \right\}$$

Glauber, 1955, 1967, 1970  
 Glauber and Matthiae, 1970  
 Bialas, Bleszynski, Czyz, 1976, ...

Optical model, high energy physics

Nuclear thickness function

Overlap function

Configuration space

Nuclear geometry (uncorrelated)

Elementary n-n interactions

Analytically  $\sim$  impossible. Nuclear short range correlations?  
 -> Monte-Carlo simulations

# SR Correlations, Gribov corrections





FIG. 2:  $\sigma_{tot}^{nA}$  vs  $p_{lab}$ . Left panel: Glauber single density approximation ( $\sigma_G$ ; dots) and Glauber plus Gribov inelastic shadowing ( $\sigma_G + \Delta\sigma_{IS}$ ; dot-dash). Right panel: Glauber ( $\sigma_G$ ; dots); Glauber plus SRC ( $\sigma_G + \sigma_{SRC}$ ; dashes); Glauber plus SRC plus Gribov inelastic shadowing ( $\sigma_G + \sigma_{SRC} + \Delta\sigma_{IS}$ ; full). Experimental data from [6, 17].

<- Avioli et al, arXiv:0708.0873




Gribov: fluctuations in the size of n decrease total nA cross-sections

Short range nucleon-nucleon correlations (SRC) + Gribov diffractive corrections are important for nA and pA collisions  
 → DIPSY detailed MC study for future accelerators


# DIPSY: Lund Dipole Cascade Model

$$\frac{dP}{dY} = \frac{\bar{\alpha}}{2\pi} d^2z \frac{(x-y)^2}{(x-z)^2(z-y)^2}, \quad \text{with } \bar{\alpha} = \frac{3\alpha_s}{\pi}.$$



The evolution of the dipole cascade in transverse coordinate space.

Based on: Mueller's dipole cascade  
Formulation of BFKL evolution  
in rapidity and transverse coordinates



A dipole-dipole interaction implies exchange of colour and reconnection

2 dipoles interact (in Born approx.)  
Multiple collisions:  
in eikonal approx., unitarity OK  
Forward amplitude and cross sections:

$$2f_{ij} = 2f(x_i, y_i | x_j, y_j) = \frac{\alpha_s^2}{4} \left[ \log \left( \frac{(x_i - y_j)^2 (y_i - x_j)^2}{(x_i - x_j)^2 (y_i - y_j)^2} \right) \right]^2.$$

Int. prob. =  $1 - e^{-2F}$ , with  $F = \sum f_{ij}$ .

$$T = 1 - e^{-F},$$


$$\sigma_{\text{inel}} = \int d^2b \langle 1 - e^{-2F(b)} \rangle = \int d^2b \langle 1 - (1 - T(b))^2 \rangle.$$

$$\sigma_{\text{el}} = \int d^2b \langle T(b) \rangle^2.$$

$$\sigma_{\text{diff}} = \int d^2b \langle T(b)^2 \rangle.$$

$$\sigma_{\text{diff ex}} = \sigma_{\text{diff}} - \sigma_{\text{el}} = \int d^2b (\langle T(b)^2 \rangle - \langle T(b) \rangle^2)$$


# DIPSY: a graphical summary




**Figure 5:** (a) A parton chain stretched between projectile and target. (b) A backbone of  $k_{\perp}$ -changing gluons in a  $(y, \ln q_{\perp}^2)$  plane. The transverse momentum of the virtual links  $k_i$  are represented by horizontal lines.

Dipole chain  
(top)

Chain split  
(bottom)




**Figure 27:** Two interactions makes a chain split in two. Dotted lines show parent structure, full lines show colour flow.



**Figure 6:** Collision of two dipole cascades in  $r$ -rapidity space. The dashed vertical line symbolizes the Lorentz frame in which the collision is viewed. The dipole splitting vertex can result in the formation of different dipole branches, and loops are formed due to multiple sub-collisions. The loop denoted by A is an effect of saturation within the cascade evolution, which can be formed via a dipole swing. Branches which do not interact, like those denoted B and C are to be treated as virtual, and to be absorbed.


Saturation  
(top)

Swing:  
rearranged  
color flow  
(bottom)



**Figure 28:** A swing between (45) and (26) causing two chains of backbone gluons to merge. Dotted lines show parent structure, full lines show colour flow. The picture is in impact-parameter space.


# Swings for glueball production?



Dipole-dipole interactions may lead to increased probability of glueball (closed color dipole loop) production  
See the talk of H. Stöcker yesterday.

# Treatment of nuclei in DIPSY

$$\rho(r) = \frac{\rho_0 \left(1 + wr^2 / R^2\right)}{1 + \exp\left((r - R) / a\right)}$$



## Electron Scattering Measurements

Nucleus	A	R	a	w
C	12	2.47	0	0
O	16	2.608	0.513	-0.051
Al	27	3.07	0.519	0
S	32	3.458	0.61	0
Ca	40	3.76	0.586	-0.161
Ni	58	4.309	0.516	-0.1308
Cu	63	4.2	0.596	0
W	186	6.51	0.535	0
Au	197	6.38	0.535	0
Pb	208	6.68	0.546	0
U	238	6.68	0.6	0

H. DeVries, C.W. De Jager, C. DeVries, 1987

$$T_A(s) = \int_{-\infty}^{+\infty} \rho_A(\vec{s}, z) dz$$

Extended Woods-Saxon charge density


Currently in DIPSY for He, O, Cu, Au and Pb

GLISSANDRO:  
(Broniowski et al)  
corrections for  
nuclear center

R(Pb,NC) = 6.40 fm  
R(Au, NC) = 6.28 fm  
R(Cu,NC) = 4.23 fm  
R(O,NC) = 2.51 fm

$$(1.1A^{1/3} - 0.656A^{-1/3}) \text{ fm}$$

# DIPSY test 1: pp cross sections




DIPSY reproduces total and elastic pp cross-sections from  $\sim 200$  GeV




# DIPSY test 2: pPb cross sections

## pPb inelastic cross sections




DIPSY reproduces  
prel. inelastic pPb  
cross-sections

# DIPSY predictions: pA




# DIPSY pA/pp ratios

pA/pp total cross section ratios



pA total cross sections




pA total cross section does not scale with A (fluctuations, swing)


DIPSY:  $\sigma_{\text{tot}}(\text{pA})$  asymptotically scales with  $(A^{1/3} + 1)^2$

# DIPSY for a future ep and eA collider

$\gamma^*p$  total cross sections



$\gamma^*O$  total cross sections




$\gamma^*p$   
DIPSY predictions


$\gamma^*A$   
DIPSY predictions

# DIPSY predictions for eA collisions

$\gamma^*O/\gamma^*p$  total cross section ratios




$\gamma^*Au/\gamma^*p$  total cross section ratios



$eO \sigma_{\text{tot}}$   
 $\sim$  scales with  $A$  :  
 fluctuations, swing  
 important


$eAu \sigma_{\text{tot}}$   
 reduced wrt  $eO$   
 but large wrt  $pp$  :  
 fluctuations, swing  
 still important

# Swing effects in DIPSY




# DIPSY predictions vs Glauber MC

Total cross sections



Total cross section ratios



DIPSY with dipole fluctuations and swing effects reduce pA cross sections by cca 5 – 15 %. Effect bigger for smaller A.

# Formalism for Black/Grey Discs

$$\sigma_{\text{tot}} = 2 \int d^2b \langle T(b) \rangle = 2\pi R^2$$

$$\sigma_{\text{el}} = \int d^2b \langle T(b) \rangle^2 = \pi R^2$$

$$\sigma_{\text{D}} = \int d^2b (\langle T(b)^2 \rangle - \langle T(b) \rangle^2) = 0$$

$$\sigma_{\text{in,ND}} = \int d^2b \langle 1 - (1 - T(b))^2 \rangle = \pi R^2.$$

In Black Disc approximation,  
3 ways to define  
the radius of the nucleons

$$\sigma_{\text{tot}}, \sigma_{\text{el}}, \sigma_{\text{in,ND}}$$

$$\sigma_{\text{tot}} = 2 \int d^2b \langle T(b) \rangle = 2\pi R^2 a$$

$$\sigma_{\text{el}} = \int d^2b \langle T(b) \rangle^2 = \pi R^2 a^2$$

$$\sigma_{\text{D}} = \int d^2b (\langle T(b)^2 \rangle - \langle T(b) \rangle^2) = \pi R^2 a(1 - a)$$

$$\sigma_{\text{in,ND}} = \int d^2b \langle 1 - (1 - T(b))^2 \rangle = \pi R^2 a.$$

In Grey Disc approx,  
R and „a”  
are usually given by

$$\sigma_{\text{tot}} \text{ and } \sigma_{\text{el}}$$

$$\sigma_{\text{el}} + \sigma_{\text{D}} = \sigma_{\text{in,ND}} = \sigma_{\text{tot}}/2$$



# DIPSY for quasi-elastic $\sigma_{\text{tot}}(\text{pPb})$


All cross-sections in barns,  
for pPb at  $\sqrt{s_{\text{NN}}} = 5 \text{ TeV}$   
Misprints fixed on 2015/06/04

Model	$\sqrt{s_{\text{NN}}}$ (TeV)	DIPSY		Black disc ( $\sigma_{\text{tot}}$ )		Black disc ( $\sigma_{\text{in}}$ )		Black disc ( $\sigma_{\text{in,ND}}$ )		Grey disc ( $\sigma_{\text{tot}}, \sigma_{\text{el}}$ )	
		5	10	5	10	5	10	5	10	5	10
$\sigma_{\text{tot}}$	(b)	3.54	3.62	3.50	3.58	3.88	3.95	3.73	3.80	3.69	3.77
$\sigma_{\text{in}}$	(b)	2.04	2.07	1.95	1.98	2.14	2.17	2.06	2.09	2.07	2.11
$\sigma_{\text{in,ND}}$	(b)	1.89	1.92	1.75	1.79	1.94	1.98	1.86	1.90	1.84	1.89
$\sigma_{\text{el}}$	(b)	1.51	1.55	1.55	1.60	1.73	1.78	1.66	1.70	1.62	1.66
$\sigma_{\text{SD,A}}$	(b)	0.085	0.086	0.198	0.192	0.204	0.198	0.200	0.195	0.083	0.085
$\sigma_{\text{SD,p}}$	(b)	0.023	0.024	-	-	-	-	-	-	-	-
$\sigma_{\text{DD}}$	(b)	0.038	0.038	-	-	-	-	-	-	0.142	0.137
$\sigma_{\text{el}*}$	(b)	1.59	1.64	1.75	1.79	1.94	1.98	1.86	1.90	1.70	1.75
$\sigma_{\text{el}*}/\sigma_{\text{in}}$		0.78	0.79	0.90	0.90	0.91	0.91	0.90	0.91	0.82	0.83
$\sigma_{\text{in,ND}}/\sigma_{\text{tot}}$		0.53	0.53	0.50	0.50	0.50	0.50	0.50	0.50	0.50	0.50

fluctuations, swings and other effects.


# What have we learned ?

Total cross section ratios



pA cross sections reduce by 10% due to dipole fluctuations + swing

Total cross section ratios



Effect is larger in eA than in pA, decreases with increasing dipole size

# What have we learned ?

Initial conditions  
for hydro evolution  
from cross-section ratios

DIPSY Monte Carlo's  
stable prediction:

$$\begin{aligned}\sigma^*/\sigma_{\text{tot}} &\sim 1/2 \\ \sigma^*/\sigma_{\text{in}} &\sim 4/5 \\ &\text{in pPb at LHC}\end{aligned}$$

Swings:  
glueballs in pPb?

eA collider:  
favourable  
as compared to pPb or dAu

# Backup slides – Questions?

# Hydro behaviour in h+p (NA22/EHS)




Fig. 11. The rapidity distributions of centrally produced pions ( $|y| < 1.5$ ) for different  $m_t$ -slices given. The curves are the fit results obtained analytically using the BL-H parameterization.





Fig. 12. The  $m_t$  distributions of centrally produced pions ( $|y| < 1.5$ ) for different  $y$ -slices given. The curves are the fit results obtained analytically using the BL-H parameterization.

Hydro behaviour in h+p reactions at  $\sqrt{s}=22 \text{ GeV}$   
 Reviewed in T. Cs, hep-ph/0001233

# Frame dependence?



DIPSY pp cross sections need to be tuned in each frame, after this step the cross section ratios are frame independent.